

Chapter 2

Genetics of Alcohol Metabolism

Vijay A. Ramchandani

Key Points

- Alcohol metabolism occurs mainly via hepatic oxidation and is governed by the catalytic properties of the alcohol-metabolizing enzymes, alcohol dehydrogenase (ADH), and aldehyde dehydrogenase (ALDH2).
- Genetic polymorphisms in *ADH1B* and *ALDH2*, and ethnic differences in the prevalence of these polymorphisms, result in increased variation in alcohol metabolism among individuals.
- Polymorphisms in *ADH1B* result in variants that code for isozymes that tend to show a faster rate of alcohol metabolism, while the *ALDH2**2 polymorphism results in a “deficient” form of *ALDH2* that causes an accumulation of acetaldehyde and its associated physiological effects.
- *ADH* and *ALDH* polymorphisms are also associated with a protective effect on the development of alcoholism. The allele frequencies of *ADH1B**2, *ADH1B**3, and *ALDH2**2 are significantly lower in individuals diagnosed with alcohol dependence compared to controls.
- Further evaluation of the factors, both genetic and environmental, regulating the rates of alcohol and acetaldehyde metabolism, will help improve our understanding of the metabolic basis and consequences of alcohol’s effects, including the risk and consequences of alcohol-related organ damage, developmental problems, as well as alcohol dependence.

Keywords Alcohol metabolism • Alcohol dehydrogenase (ADH) • Aldehyde dehydrogenase (ALDH) • Genetic polymorphism • Ethnic differences • Cytochrome P450 • Catalase • Pharmacogenetics

Introduction

Ethanol (also referred to as alcohol in this chapter) is probably the most widely investigated drug in the world, not only because of its ubiquitous use and its widespread abuse but also because of its unique pharmacological properties. Following administration, systemic concentrations of alcohol are a consequence of the absorption, distribution, and metabolism of alcohol, which display very unique characteristics and demonstrate substantial interindividual variability [1]. As the pharmacological

V.A. Ramchandani, Ph.D. (✉)

Laboratory of Clinical and Translational Studies, National Institute on Alcohol Abuse and Alcoholism,
National Institutes of Health, 10 Center Drive, MSC 1540, Bldg 10 CRC, Rm 2-2352,
Bethesda, MD 20892-1540, USA
e-mail: vijayr@mail.nih.gov

effects of alcohol depend on its systemic concentrations, variability in the pharmacokinetics of alcohol can have a significant impact on its pharmacodynamic effects.

Following oral ingestion, alcohol is absorbed by passive diffusion, primarily from the small intestine [2, 3]. The rate of absorption depends on several factors, both genetic and environmental, and is highly variable. Some of these factors include the volume, concentration, and nature of the alcoholic beverage [2, 4, 5]; the rate of drinking [4]; the fed or fasted state [6]; the nature and composition of food [6, 7]; the rate of gastric emptying [8, 9]; the gender differences in first-pass metabolism [10, 11]; and other drugs including histamine (H1) receptor antagonists like cimetidine and ranitidine [12, 13]. Ethanol is a small polar molecule and its volume of distribution is comparable to total body water [3]. No plasma protein binding has been reported for alcohol. Elimination of alcohol occurs primarily through metabolism with small fractions of the administered dose being excreted in the breath (0.7%), sweat (0.1%), and urine (0.3%) [3]. Alcohol metabolism occurs mainly via hepatic oxidation and is governed by the catalytic properties of the alcohol-metabolizing enzymes, alcohol dehydrogenase (ADH) and aldehyde dehydrogenase (ALDH). The cytochrome P450 enzymes (CYP2E1) and catalase also contribute to alcohol metabolism and alcohol-related cytotoxicity under specific circumstances [14].

Alcohol metabolic rates show a considerable degree of interindividual and ethnic variability, in part due to allelic variants of the genes encoding ADH and ALDH producing functionally different isozymes [15–17]. Functional polymorphisms of the *ADH1B* and *ALDH2* genes have been shown to increase the variance in alcohol metabolism among individuals. Additionally, a multitude of environmental factors can influence the metabolic regulation of alcohol metabolism, which results in a large three- to four-fold variance in the alcohol elimination rate in humans [16, 18]. Factors that have been shown to be important determinants of alcohol metabolism include age [19, 20], gender [21, 22], ethnicity and genetics [21, 23–26], body mass and liver size [22], as well as environmental factors such as food intake [27].

This chapter will focus on genetic variation in the alcohol-metabolizing enzymes and its impact on the metabolism of alcohol.

Alcohol-Metabolizing Enzymes and Genetic Aspects

Alcohol Dehydrogenase

The genes for the human *ADH* family cluster in a region of chromosome 4q21 spanning ~370 kb [28]. The alcohol dehydrogenase (*ADH*) gene family encodes oxidative enzymes that metabolize a wide variety of alcohols including ethanol, retinol, other aliphatic alcohols, hydroxysteroids, and lipid peroxidation products [15, 17]. Currently, seven human *ADH* genes have been identified and organized into five classes based on amino acid sequence alignments, catalytic properties, and patterns of tissue-specific expression [29]. Human ADH enzyme is a dimeric molecule, arising from the association of different subunits expressed by the seven genes. Thus, there are over 20 ADH isozymes that vary greatly with regard to the types of alcohols they preferentially metabolize and the maximal rate at which they oxidize ethanol [15]. The five classes of ADH are divided according to their subunit and isozyme composition (Table 2.1).

The class I isozymes are found in liver and consist of homo- and heterodimeric forms of the three subunits (i.e., $\alpha\alpha$, $\alpha\beta$, $\beta\beta$, $\beta\gamma$, $\gamma\gamma$, etc.). Classes II, III, and IV enzymes are homodimeric forms of the π , χ , and σ subunits, respectively. All the class I ADHs metabolize ethanol and are inhibited by pyrazole derivatives [17]. The *ADH1* subunits share about 94% sequence identity. The relative order of catalytic efficiency (kcat/Km) for ethanol oxidation at ethanol concentrations of about 100 mg% and saturating coenzyme NAD⁺ concentration (0.5 mM) is $\beta2 > \beta1 > \gamma1 > \gamma2 \approx \sigma >> \beta3 > \alpha >> \pi$. However, the relative order of kcat at saturating concentrations of both ethanol and NAD⁺ is $\sigma > \beta3 \approx \beta2 > \gamma1 > \gamma2 \approx \pi > \beta1$. Thus, the relative contributions of each of the ADH isozymes to ethanol oxidation change with the hepatic concentration of alcohol [16, 17, 30].

Table 2.1 Nomenclature for alcohol dehydrogenase genes

ADH class	Official gene nomenclature	Former gene nomenclature	Enzyme subunit nomenclature	Km for ethanol [mM]
I	<i>ADH1A</i>	<i>ADH1</i>	α	4.0
I	<i>ADH1B*1</i>	<i>ADH2*1</i>	β 1	0.05
I	<i>ADH1B*2</i>	<i>ADH2*2</i>	β 2	0.9
I	<i>ADH1B*3</i>	<i>ADH2*3</i>	β 3	40
I	<i>ADH1C*1</i>	<i>ADH3*1</i>	γ 1	1.0
I	<i>ADH1C*2</i>	<i>ADH3*2</i>	γ 2	6.0
II	<i>ADH4</i>	<i>ADH4</i>	π	30
III	<i>ADH5</i>	<i>ADH5</i>	χ	>1,000
IV	<i>ADH7</i>	<i>ADH7</i>	σ	30
V	<i>ADH6</i>	<i>ADH6</i>	Not identified	?

For official gene nomenclature, go to: <http://www.genenames.org/genefamilies/ADH> (HUGO Gene Nomenclature Committee at the European Bioinformatics Institute)

The human *ADH* genes are differentially expressed in different tissues, and this is a fundamental determinant of the physiological consequences of alcohol metabolism in specific cells and tissues [31]. The liver contains a large amount of ADH (about 3% of soluble protein) and expresses the widest number of different isozymes. ADH4 (π -ADH) is solely expressed in liver. Only ADH7 (σ -ADH) is not highly expressed in liver. ADH5 (χ -ADH) is ubiquitously expressed in human tissues. *ADH1C*, *ADH4*, *ADH5*, and *ADH7* are expressed in gastrointestinal tissues. The expression of ADH6 in humans and its role in ethanol metabolism remains to be elucidated. The expression of ADH in other tissues such as skeletal muscle, and the quantitative significance of muscle ADH metabolism (because of the large proportion of muscle mass in the body), also remains to be determined.

In addition to ethanol, alcohol dehydrogenases also oxidize several “physiological” alcohols with high catalytic efficiency including retinol, ω -hydroxy fatty acids, hydroxysteroids, and hydroxy derivatives of dopamine and epinephrine metabolites [30, 32]. Oxidation of these alcohols can be inhibited by ethanol, and therefore the role of ethanol substrate competition is an important issue in alcohol-related toxicology. Another important issue is the regional expression of ADHs in brain and their potential role in the local formation of acetaldehyde, which may be centrally active, possessing stimulant as well as sedative/hypnotic effects [33–35].

Genetic Variation

Single nucleotide polymorphisms (SNP) have been identified at the *ADH1B* and *ADH1C* loci [15, 17, 31]. Variant alleles of *ADH1B* result in the β 1, β 2, and β 3 subunits, while variants in *ADH1C* result in the γ 1 and γ 2 subunits. The resulting subunits have different catalytic activities for ethanol (see Table 2.1). Additionally, the *ADH1B* alleles appear with different frequencies in different racial groups, with the *ADH1B*1* form predominating in Caucasian and African-descent populations, and *ADH1C*2* predominating in East Asian populations (e.g., Chinese and Japanese), and also found in about 25% of Caucasians with Jewish ancestry. The *ADH1B*3* form is found in about 25% of individuals of African descent. With respect to the *ADH1C* polymorphism, *ADH1C*1* and *ADH1C*2* appear with about equal frequency in Caucasians, but *ADH1C*1* predominates in African-descent and East Asian populations [36]. Recently, a novel polymorphism was identified in *ADH1C*. This polymorphism results in an allele that codes for a subunit with a proline to threonine substitution in position 351 and has been described in Native Americans [37]. However, the catalytic activity of the isozyme coded by this variant and its effect on the overall elimination of alcohol remains to be determined.

There are additional SNPs that have been identified in the noncoding regions of the *ADH* genes. Several of these SNPs have been shown to affect the expression of *ADH* genes [31, 38] and may be

associated with alcoholism risk [39]; however, the effect of these variations on the catalytic activity of ADH and effect on the overall metabolism of alcohol remains to be established.

Aldehyde Dehydrogenase

Acetaldehyde is the first metabolic product of ethanol metabolism and is itself metabolized via oxidation by the NAD⁺-dependent aldehyde dehydrogenase (ALDH). Several isozymes of ALDH, differing in kinetic properties and tissue distribution, have been detected in human organs and tissues [15]. Currently, 19 putatively functional *ALDH* genes have been identified in the human genome [40, 41]. However, only the *ALDH1* (*ALDH1A1*) and *ALDH2* genes encode the class I and class II isozymes that are involved in acetaldehyde oxidation. *ALDH1* is the cytosolic form distributed ubiquitously in tissues including brain. It exhibits relatively low catalytic activity ($K_m \sim 30 \mu\text{M}$) for acetaldehyde oxidation. *ALDH2* is the mitochondrial enzyme that is highly expressed in liver and stomach [42]. It exhibits high catalytic activity ($K_m \sim 3 \mu\text{M}$) for acetaldehyde oxidation and is primarily responsible for acetaldehyde oxidation *in vivo*.

Genetic Variation

The best-known genetic polymorphism in *ALDH* genes is in *ALDH2*. The allelic variants are *ALDH2*1* and *ALDH2*2*, encoding for the high-activity and low-activity forms of the subunits respectively. The low-activity form arises from a single amino acid exchange (glutamine to lysine substitution at position 487) at the coenzyme-binding site of the enzyme subunit [15, 17]. This results in a 100-fold increase in the K_m for acetaldehyde [43]. This very prominent variant allele has been seen in about half of the East Asian populations studied (including the Han Chinese, Taiwanese, and Japanese) [44, 45]. It has not been observed in populations of Caucasian origin. It exhibits virtually no acetaldehyde oxidizing activity *in vitro* and represents the “deficient” phenotype seen in these Asian populations [46]. Individuals who are heterozygous or homozygous for *ALDH2*2* show accumulation of acetaldehyde levels and the characteristic sensitivity reaction (facial flushing, increased skin temperature and heart rate) following alcohol intake [26, 28, 47, 48].

Cytochrome P450 Enzymes

A small fraction of an ingested dose of ethanol is metabolized by enzymes other than ADH. Metabolism of ethanol by the so-called microsomal ethanol oxidizing system (MEOS) accounts for the major non-ADH system [14, 49]. MEOS consists primarily of the cytochrome P450 isoform, P4502E1 (*CYP2E1*), along with other P450 enzymes, and is the major alternative system that catalyzes the NADPH- and O₂-dependent oxidation of ethanol to form acetaldehyde, NADP⁺, and water. Like other cytochrome P450 enzymes, the primary role of *CYP3E1* is the metabolism of alcohol and other xenobiotics. While *CYP2E1* accounts for a much smaller fraction of ethanol oxidation than the ADH system under normal conditions, it represents a major adaptive response of alcohol metabolism with chronic ethanol consumption [49]. This is due to the direct effect of chronic ethanol consumption on the expression of hepatic *CYP2E1*. In humans, there is an induction of *CYP2E1* with chronic alcohol consumption that can be followed by a decrease in activity associated with generalized hepatic injury and loss of function. There are two mechanisms postulated for *CYP2E1* induction: (1) a posttranslational mechanism involving mRNA stabilization and protection of the expressed protein against degradation and (2) a

direct transcriptional regulation of *CYP2E1* expression, generally following high exposures to ethanol. The expression of *CYP2E1* is influenced by factors such as diet (lipids, carbohydrates) and hormones (thyroid hormones, glucocorticoids, steroids, pituitary hormones). The induction of *CYP2E1* may result in higher levels of toxic metabolites of other xenobiotics as well as the generation of superoxide radicals, which may contribute to the increased risk of alcohol-related liver disease as well as cancer.

Genetic Variation

A number of different *CYP2E1* polymorphisms have been identified [15, 50]. A variant allele called *5B has been identified in the 5'-flanking region of the *CYP2E1* gene. This allele has been shown to be differentially expressed in different racial populations, and the variant allele (previously labeled as the *c2* allele) has been found to be associated with higher transcriptional activity, protein levels, and enzyme activity than the common wild-type *c1* allele [51]. The influence of this polymorphism on alcohol elimination was examined in one study in Japanese alcoholics and control and indicated that the presence of the *c2* allele (heterozygous or homozygous) may be associated with higher alcohol metabolic rates but only at blood alcohol levels greater than 0.25% (g/dL) [52]. Studies have identified additional genetic variation that may be relevant to alcohol, including the *1D allele, which has been found at higher frequency in Chinese (23%) and African-Americans (31%) than in Caucasians (1–7%) [53, 54]. Studies in African-Americans have further shown higher levels of *CYP2E1* inducibility following alcohol intake as measured by oxidation of the *CYP2E1* substrate chlorzoxazone. However, the influence of this polymorphism on alcohol metabolism remains to be determined. Much work needs to be done to understand mechanisms for transcriptional and posttranslational regulation of the *CYP2E1* genes and their role in alcohol metabolism and alcohol-related liver disease [49].

Catalase

Catalase is an enzyme that catalyzes the hydrogen peroxide (H₂O₂)-dependent oxidation of ethanol yielding acetaldehyde and two molecules of water. It is found in the cytosol and mitochondria but its main expression and function is in peroxisomes. Most studies indicate that it contributes very little to total ethanol elimination because of the limited availability of hydrogen peroxide [14, 55]. However, the activation of peroxisomal catalase by increased generation of hydrogen peroxide via peroxisomal β-oxidation can lead to a hypermetabolic state and a swift increase in alcohol metabolism under some conditions [56]. This state may contribute to alcohol-related inflammation and necrosis in alcoholic liver disease. Additional studies have suggested that catalase may be involved in the metabolism of alcohol to acetaldehyde in the brain. This has led to implications of a role for acetaldehyde in mediating some of the behavioral effects of alcohol [35]. However, further research is needed to clarify the pharmacokinetics and central pharmacodynamic effects of acetaldehyde and its role in the pharmacology of alcohol.

ADH and ALDH Polymorphisms: Influence on Alcohol Metabolism

Functional polymorphisms of genes for the alcohol-metabolizing enzymes *ADH* and *ALDH2*, and differences in the prevalence of the polymorphic alleles in different ethnic populations, have resulted in several studies examining ethnic differences in alcohol metabolism and the influence of *ADH1B*, *ADH1C*, and *ALDH2* genotypes. The isozymes encoded by the polymorphic alleles have very different catalytic properties in vitro, as described earlier in this chapter, and would be expected to exert influences on an individual's alcohol metabolic rate.

One of the first studies examining the influence of *ADH* and *ALDH* polymorphisms on alcohol metabolism was done by Mizoi et al. [23] in 68 healthy Japanese subjects. Subjects were genotyped for *ADH1B* as well as *ALDH2* polymorphisms and alcohol disappearance rates (mg/ml/h), and elimination rates (mg/kg/h) were compared among the groups classified based on genotypes of both *ADH1B* (*ADH1B**1/*1, *ADH1B**1/*2, and *ADH1B**2/*2) and *ALDH2*. Results indicated that there were no differences in alcohol metabolism among the *ADH1B* genotypes; however, there were marked differences among the *ALDH2* genotypes with regard to alcohol metabolism. Other studies in Asians have also failed to demonstrate an effect of the *ADH1B**2 allele on alcohol metabolism after controlling for the *ALDH2**2 polymorphism. This is discussed further below.

Studies in Jewish individuals possessing the *ADH1B**2 polymorphism have provided a clearer picture of the effect of this variant on alcohol metabolism, Neumark et al. [57] conducted a study in young healthy Jewish males to assess the effect of the *ADH1B* polymorphism on alcohol elimination rates measured using the alcohol clamp [58]. Results revealed a significantly higher alcohol elimination rates in subjects carrying the *ADH1B**2 allele (heterozygotes and homozygotes) compared with *ADH1B**1 homozygotes [57, 59]. As the Jewish do not show polymorphisms of the *ALDH2* genes, this appears to be a direct effect of *ADH1B* genotypes on alcohol metabolism.

Thomasson et al. [21] examined the influence of the *ADH1B**3 polymorphism on alcohol metabolism in a sample of 112 African-American subjects, selected by genotype. In this study, subjects received an oral dose of alcohol and alcohol disappearance rates were determined from the slope of the pseudo-linear portion of the blood ethanol concentration vs. time curves. Results revealed that subjects carrying the *ADH1B**3 allele (heterozygotes and homozygotes) showed a higher alcohol disappearance rate (mg% per h) for compared to *ADH1B**1 homozygotes. A more recent study in African-Americans failed to demonstrate an effect of the *ADH1B**3 polymorphism on breath alcohol concentrations following a moderate oral dose of alcohol in 91 African-Americans [60]. A study in Native Americans also showed that subjects with *ADH1B**3 alleles had a trend toward higher alcohol elimination rates than subjects with *ADH1B**1 [24]. However, this difference was not statistically significant probably because of the small number of subjects possessing the *ADH1B**3 genotype in the study and the low frequency of occurrence of this genotype (~7%) in this ethnic group. Earlier studies in Native Americans have previously demonstrated higher alcohol elimination rates compared to those reported in Caucasians; however, *ADH* genotypes were not determined in these studies [61, 62].

The influence of *ALDH2* polymorphisms on alcohol metabolism has been studied more extensively, although almost exclusively in Asian subjects, mainly because of the high frequency of the polymorphism in this population. Most of these studies have compared peak concentrations of alcohol and acetaldehyde as well as peak responses on subjective and cardiovascular measures and flushing across *ADH1B* and *ALDH2* genotypes, with generally consistent results. In general, individuals who are heterozygous or homozygous for *ALDH2**2 show increased acetaldehyde levels following alcohol administration [23, 25, 28, 47, 63–65]. Some studies have also demonstrated significant increases in ethanol concentrations and area under the ethanol concentration time curves [63, 65], possibly due to product inhibition of the ADH activity by acetaldehyde. However, other studies have shown accumulation of acetaldehyde in subjects carrying the *ALDH2**2 allele without any difference in alcohol concentrations or elimination rates [25, 26].

Given the high frequency of the *ADH1B**2 and *ALDH2**2 alleles in Asians, it is important to understand the contribution of each polymorphism to the observed differences in blood alcohol and acetaldehyde levels following alcohol administration. There are only a few studies that have actually estimated and compared alcohol disappearance rates or elimination rates among *ADH1B* and/or *ALDH2* genotypes. In the study by Mizoi et al. [23] described above, peak acetaldehyde levels, alcohol disappearance rates (mg/ml/h), and elimination rates (mg/kg/h) were compared among subjects classified into groups based on genotypes of both *ADH1B* and *ALDH2* (*ALDH2**1/*1, *ALDH2**1/*2, and *ALDH2**2/*2). Results indicated that subjects homozygous for *ALDH2**1/*1 showed no increase in acetaldehyde levels regardless of their *ADH1B* genotype. There was a progressive increase in peak acetaldehyde levels in subjects with the *ALDH2**1/*2 and *ALDH2**2/*2 genotypes. Both alcohol

disappearance rates and elimination rates showed significant differences among the *ALDH2* genotypes and decreased in the following order: *ALDH2*1/*1* > *ALDH2*1/*2* > *ALDH2*2/*2*. A study in Chinese men indicated that the presence of the *ALDH2*2* allele was associated with slower alcohol metabolism following oral administration, while in individuals homozygous for *ALDH2*1*, the presence of two *ADH2*2* alleles correlated with slightly faster alcohol metabolism [66]. Studies by Peng et al. [26, 48, 63] have demonstrated a clear effect of *ALDH2* genotype on alcohol and acetaldehyde metabolism, as well as the lack of significant effect of *ADH1B* polymorphism on acetaldehyde metabolism. In fact, most studies in Asians have not demonstrated that the *ADH1B*2* allele is associated with differences in alcohol metabolism after controlling for the *ALDH2* [25, 47, 67].

A recent effort in understanding the influence of genetic variation in alcohol-metabolizing enzymes on alcohol metabolism has focused on the use of association analysis in a large cohort of twin pairs of Caucasian ancestry. In these studies, 103 SNPs spanning the *ADH* gene family were examined for association with measures of alcohol metabolism following oral alcohol challenge in this sample. Results indicated significant associations between alcohol elimination rates and *ADH1A*, *ADH1B*, *ADH1C*, as well as *ADH7* genes [68, 69]. These studies point to a role for *ADH7* in the metabolism of alcohol; however, more work is needed to clarify the influence of this isoform, and its associated genetic variation, on alcohol elimination rates in humans.

In summary, genetic polymorphisms of *ADH* and *ALDH* result in alterations in the metabolism of alcohol and/or acetaldehyde. Polymorphisms in *ADH1B* result in variants that code for isozymes that tend to show a faster rate of alcohol metabolism, while the *ALDH2*2* polymorphism results in a “deficient” form of *ALDH2* that causes an accumulation of acetaldehyde and its associated physiological effects.

***ADH* and *ALDH* Polymorphisms: Association with Alcohol Dependence**

Functional polymorphisms of the alcohol-metabolizing enzymes *ADH* and *ALDH2* can also exert important effects on the biological effects of alcohol [26, 70]. In fact, the *ADH* and *ALDH* genes are the only genes which have been firmly established to influence vulnerability to alcohol dependence or alcoholism [17, 36]. Studies have demonstrated unequivocally that the allele frequencies of *ADH1B*2*, *ADH1B*3*, and *ALDH2*2* are significantly decreased in subjects diagnosed with alcohol dependence as compared with the general population of East Asians, including the Japanese, Han Chinese, and Koreans [39, 44, 45, 67, 71–76]. The *ALDH2*2* allele and the *ADH1B*2* allele also significantly influence drinking behavior in nonalcoholic individuals. Association between reduced alcohol consumption or reduced risk of alcohol dependence and the *ADH1B*2* variant allele has recently been found in other ethnic groups that do not carry the *ALDH2*2* allele, including Europeans [77–80], Jews in Israel [81, 82], as well as Mongolians in China [45], and the Atayal natives of Taiwan [83]. Recent studies have also shown a protective association between the *ADH1B*3* allele and alcohol dependence in Native Americans. [84, 85] Finally, studies have indicated that the *ADH1B*3* allele may be protective against alcohol-related problems in infants born to African-American mothers who may have consumed alcohol during pregnancy [86–89].

Summary

There has been substantial progress in the field of alcohol pharmacogenetics to characterize differences in alcohol metabolism in subjects exhibiting polymorphic genotypes of the alcohol-metabolizing enzymes. The impact of functional variation in *ADH1B* and *ALDH2* genes on alcohol metabolism have been fairly well characterized; however, there are large interindividual differences in alcohol elimination

rates that still remain unexplained. Of potential significance in this regard may be polymorphisms in *ADH4* [90, 91], *ADH7* [39, 69], and *ALDH1A1* [92, 93] as well as the promoter regions of *ALDH2* [94]. Further studies are needed to evaluate the influence of these polymorphisms on the activity of ADH and ALDH and on alcohol levels and elimination rates in individuals, as well as on the physiological response to alcohol consumption and alcoholism. Recent integrated approaches examining the associations of *ADH* and *ALDH2* gene variation with alcohol metabolism, response, drinking behavior, and alcohol dependence in large samples [78] might be particularly useful in this regard.

Studies in monozygotic and dizygotic twins have shown that the heritability (i.e., genetic component of variance) of alcohol metabolic rates is about 50% [95, 96]. Further evaluation of the factors, both genetic and environmental, regulating the rates of alcohol and acetaldehyde metabolism, will help improve our understanding of the metabolic basis and consequences of alcohol's effects, including the risk and consequences of alcohol-related organ damage, developmental problems, as well as alcohol dependence.

Acknowledgements Division of Intramural Clinical and Biological Research, National Institute on Alcohol Abuse and Alcoholism, National Institutes of Health.

References

1. Norberg A, Jones AW, Hahn RG, Gabrielsson JL. Role of variability in explaining ethanol pharmacokinetics: research and forensic applications. *Clin Pharmacokinet.* 2003;42:1–31.
2. Wilkinson PK, Sedman AJ, Sakmar E, Kay DR, Wagner JG. Pharmacokinetics of ethanol after oral administration in the fasting state. *J Pharmacokinet Biopharm.* 1977;5:207–24.
3. Holford NH. Clinical pharmacokinetics of ethanol. *Clin Pharmacokinet.* 1987;13:273–92.
4. O'Neill B, Williams AF, Dubowski KM. Variability in blood alcohol concentrations. Implications for estimating individual results. *J Stud Alcohol.* 1983;44:222–30.
5. Dubowski KM. Absorption, distribution and elimination of alcohol: highway safety aspects. *J Stud Alcohol Suppl.* 1985;10:98–108.
6. Sedman AJ, Wilkinson PK, Sakmar E, Weidler DJ, Wagner JG. Food effects on absorption and metabolism of alcohol. *J Stud Alcohol.* 1976;37:1197–214.
7. Jones AW, Jonsson KA, Kechagias S. Effect of high-fat, high-protein, and high-carbohydrate meals on the pharmacokinetics of a small dose of ethanol. *Br J Clin Pharmacol.* 1997;44:521–6.
8. Mushambi MC, Bailey SM, Trotter TN, Chadd GD, Rowbotham DJ. Effect of alcohol on gastric emptying in volunteers. *Br J Anaesth.* 1993;71:674–6.
9. Kalant H. Effects of food and body composition on blood alcohol curves. *Alcohol Clin Exp Res.* 2000;24:413–4.
10. Frezza M, di Padova C, Pozzato G, Terpin M, Baraona E, Lieber CS. High blood alcohol levels in women. The role of decreased gastric alcohol dehydrogenase activity and first-pass metabolism. *N Engl J Med.* 1990;322:95–99.
11. Ammon E, Schafer C, Hofmann U, Klotz U. Disposition and first-pass metabolism of ethanol in humans: is it gastric or hepatic and does it depend on gender? *Clin Pharmacol Ther.* 1996;59:503–13.
12. Gupta AM, Baraona E, Lieber CS. Significant increase of blood alcohol by cimetidine after repetitive drinking of small alcohol doses. *Alcohol Clin Exp Res.* 1995;19:1083–7.
13. Arora S, Baraona E, Lieber CS. Alcohol levels are increased in social drinkers receiving ranitidine. *Am J Gastroenterol.* 2000;95:208–13.
14. Lieber CS. Microsomal ethanol-oxidizing system (MEOS): the first 30 years (1968–1998)—a review. *Alcohol Clin Exp Res.* 1999;23:991–1007.
15. Agarwal DP. Genetic polymorphisms of alcohol metabolizing enzymes. *Pathol Biol (Paris).* 2001;49:703–9.
16. Ramchandani VA, Bosron WF, Li TK. Research advances in ethanol metabolism. *Pathol Biol (Paris).* 2001;49:676–82.
17. Hurlley TD, Edenberg HJ, Li TK. Pharmacogenetics of alcoholism. In: Licinio J, Wong ML, editors. *Pharmacogenomics: the search for individualized therapeutics.* Weinheim: Wiley-VCH; 2002. p. 417–41.
18. Eckardt MJ, File SE, Gessa GL, et al. Effects of moderate alcohol consumption on the central nervous system. *Alcohol Clin Exp Res.* 1998;22:998–1040.
19. Vestal RE, McGuire EA, Tobin JD, Andres R, Norris AH, Mezey E. Aging and ethanol metabolism. *Clin Pharmacol Ther.* 1977;21:343–54.

20. Jones AW, Neri A. Age-related differences in blood ethanol parameters and subjective feelings of intoxication in healthy men. *Alcohol Alcohol*. 1985;20:45–52.
21. Thomasson HR, Beard JD, Li TK. ADH2 gene polymorphisms are determinants of alcohol pharmacokinetics. *Alcohol Clin Exp Res*. 1995;19:1494–9.
22. Kwo PY, Ramchandani VA, O'Connor S, et al. Gender differences in alcohol metabolism: relationship to liver volume and effect of adjusting for body mass. *Gastroenterology*. 1998;115:1552–7.
23. Mizoi Y, Yamamoto K, Ueno Y, Fukunaga T, Harada S. Involvement of genetic polymorphism of alcohol and aldehyde dehydrogenases in individual variation of alcohol metabolism. *Alcohol Alcohol*. 1994;29:707–10.
24. Wall TL, Garcia-Andrade C, Thomasson HR, Cole M, Ehlers CL. Alcohol elimination in Native American Mission Indians: an investigation of interindividual variation. *Alcohol Clin Exp Res*. 1996;20:1159–64.
25. Wall TL, Peterson CM, Peterson KP, et al. Alcohol metabolism in Asian-American men with genetic polymorphisms of aldehyde dehydrogenase. *Ann Intern Med*. 1997;127:376–9.
26. Peng GS, Yin JH, Wang MF, Lee JT, Hsu YD, Yin SJ. Alcohol sensitivity in Taiwanese men with different alcohol and aldehyde dehydrogenase genotypes. *J Formos Med Assoc*. 2002;101:769–74.
27. Ramchandani VA, Kwo PY, Li TK. Effect of food and food composition on alcohol elimination rates in healthy men and women. *J Clin Pharmacol*. 2001;41:1345–50.
28. Peng GS, Yin SJ. Effect of the allelic variants of aldehyde dehydrogenase *ALDH2**2 and alcohol dehydrogenase *ADH1B**2 on blood acetaldehyde concentrations. *Hum Genomics*. 2009;3:121–7.
29. Duester G, Farres J, Felder MR, et al. Recommended nomenclature for the vertebrate alcohol dehydrogenase gene family. *Biochem Pharmacol*. 1999;58:389–95.
30. Edenberg HJ, Bosron WF. Alcohol dehydrogenases. In: Guengerich FP, editor. *Biotransformation*. New York: Pergamon; 1997. p. 119–31.
31. Edenberg HJ. Regulation of the mammalian alcohol dehydrogenase genes. *Prog Nucleic Acid Res Mol Biol*. 2000;64:295–341.
32. Boleda MD, Saubi N, Farres J, Pares X. Physiological substrates for rat alcohol dehydrogenase classes: aldehydes of lipid peroxidation, omega-hydroxyfatty acids, and retinoids. *Arch Biochem Biophys*. 1993;307:85–90.
33. Hunt WA. Role of acetaldehyde in the actions of ethanol on the brain – a review. *Alcohol*. 1996;13:147–51.
34. Zimatkin SM, Liopo AV, Deitrich RA. Distribution and kinetics of ethanol metabolism in rat brain. *Alcohol Clin Exp Res*. 1998;22:1623–7.
35. McBride WJ, Li TK, Deitrich RA, Zimatkin S, Smith BR, Rodd-Henricks ZA. Involvement of acetaldehyde in alcohol addiction. *Alcohol Clin Exp Res*. 2002;26:114–9.
36. Li TK. Pharmacogenetics of responses to alcohol and genes that influence alcohol drinking. *J Stud Alcohol*. 2000;61:5–12.
37. Osier MV, Pakstis AJ, Goldman D, Edenberg HJ, Kidd JR, Kidd KK. A proline-threonine substitution in codon 351 of *ADH1C* is common in Native Americans. *Alcohol Clin Exp Res*. 2002;26:1759–63.
38. Chen HJ, Tian H, Edenberg HJ. Natural haplotypes in the regulatory sequences affect human alcohol dehydrogenase 1 C (*ADH1C*) gene expression. *Hum Mutat*. 2005;25:150–5.
39. Edenberg HJ, Xuei X, Chen HJ, et al. Association of alcohol dehydrogenase genes with alcohol dependence: a comprehensive analysis. *Hum Mol Genet*. 2006;15:1539–49.
40. Sophos NA, Vasiliou V. Aldehyde dehydrogenase gene superfamily: the 2002 update. *Chem Biol Interact*. 2003;143–144:5–22.
41. Vasiliou V, Nebert DW. Analysis and update of the human aldehyde dehydrogenase (ALDH) gene family. *Hum Genomics*. 2005;2:138–43.
42. Yoshida A, Rzhetsky A, Hsu LC, Chang C. Human aldehyde dehydrogenase gene family. *Eur J Biochem*. 1998;251:549–57.
43. Steinmetz CG, Xie P, Weiner H, Hurley TD. Structure of mitochondrial aldehyde dehydrogenase: the genetic component of ethanol aversion. *Structure*. 1997;5:701–11.
44. Thomasson HR, Edenberg HJ, Crabb DW, et al. Alcohol and aldehyde dehydrogenase genotypes and alcoholism in Chinese men. *Am J Hum Genet*. 1991;48:677–81.
45. Shen YC, Fan JH, Edenberg HJ, et al. Polymorphism of ADH and ALDH genes among four ethnic groups in China and effects upon the risk for alcoholism. *Alcohol Clin Exp Res*. 1997;21:1272–7.
46. Crabb DW, Edenberg HJ, Bosron WF, Li TK. Genotypes for aldehyde dehydrogenase deficiency and alcohol sensitivity The inactive ALDH2(2) allele is dominant. *J Clin Invest*. 1989;83:314–6.
47. Chen YC, Peng GS, Wang MF, Tsao TP, Yin SJ. Polymorphism of ethanol-metabolism genes and alcoholism: correlation of allelic variations with the pharmacokinetic and pharmacodynamic consequences. *Chem Biol Interact*. 2009;178:2–7.
48. Peng GS, Chen YC, Tsao TP, Wang MF, Yin SJ. Pharmacokinetic and pharmacodynamic basis for partial protection against alcoholism in Asians, heterozygous for the variant ALDH2*2 gene allele. *Pharmacogenet Genomics*. 2007;17:845–55.
49. Lieber CS. The discovery of the microsomal ethanol oxidizing system and its physiologic and pathologic role. *Drug Metab Rev*. 2004;36:511–29.

50. Neafsey P, Ginsberg G, Hattis D, Johns DO, Guyton KZ, Sonawane B. Genetic polymorphism in CYP2E1: population distribution of CYP2E1 activity. *J Toxicol Environ Health B Crit Rev*. 2009;12:362–88.
51. Hayashi S, Watanabe J, Kawajiri K. Genetic polymorphisms in the 5'-flanking region change transcriptional regulation of the human cytochrome P450IIE1 gene. *J Biochem*. 1991;110:559–65.
52. Ueno Y, Adachi J, Imamichi H, Nishimura A, Tatsuno Y. Effect of the cytochrome P-450IIE1 genotype on ethanol elimination rate in alcoholics and control subjects. *Alcohol Clin Exp Res*. 1996;20:17A–21.
53. Hu Y, Hakkola J, Oscarson M, Ingelman-Sundberg M. Structural and functional characterization of the 5'-flanking region of the rat and human cytochrome P450 2E1 genes: identification of a polymorphic repeat in the human gene. *Biochem Biophys Res Commun*. 1999;263:286–93.
54. McCarver DG, Byun R, Hines RN, Hichme M, Wegenek W. A genetic polymorphism in the regulatory sequences of human CYP2E1: association with increased chlorzoxazone hydroxylation in the presence of obesity and ethanol intake. *Toxicol Appl Pharmacol*. 1998;152:276–81.
55. Crabb DW. Ethanol oxidizing enzymes: roles in alcohol metabolism and alcoholic liver disease. *Prog Liver Dis*. 1995;13:151–72.
56. Bradford BU, Enomoto N, Ikejima K, et al. Peroxisomes are involved in the swift increase in alcohol metabolism. *J Pharmacol Exp Ther*. 1999;288:254–9.
57. Neumark YD, Friedlander Y, Durst R, et al. Alcohol dehydrogenase polymorphisms influence alcohol-elimination rates in a male Jewish population. *Alcohol Clin Exp Res*. 2004;28:10–4.
58. Ramchandani VA, O'Connor S. Studying alcohol elimination using the alcohol clamp method. *Alcohol Res Health*. 2006;29:286–90.
59. Ramchandani VA, O'Connor S, Neumark Y, Zimmermann US, Morzorati SL, de Wit H. The alcohol clamp: applications, challenges, and new directions—an RSA 2004 symposium summary. *Alcohol Clin Exp Res*. 2006;30:155–64.
60. McCarthy DM, Pedersen SL, Lobos EA, Todd RD, Wall TL. *ADH1B*3* and response to alcohol in African-Americans. *Alcohol Clin Exp Res*. 2010;34:1274–81.
61. Farris JJ, Jones BM. Ethanol metabolism in male American Indians and whites. *Alcohol Clin Exp Res*. 1978;2:77–81.
62. Reed TE, Kalant H, Gibbins RJ, Kapur BM, Rankin JG. Alcohol and acetaldehyde metabolism in Caucasians, Chinese and Amerinds. *Can Med Assoc J*. 1976;115:851–5.
63. Peng GS, Wang MF, Chen CY, et al. Involvement of acetaldehyde for full protection against alcoholism by homozygosity of the variant allele of mitochondrial aldehyde dehydrogenase gene in Asians. *Pharmacogenetics*. 1999;9:463–76.
64. Enomoto N, Takase S, Yasuhara M, Takada A. Acetaldehyde metabolism in different aldehyde dehydrogenase-2 genotypes. *Alcohol Clin Exp Res*. 1991;15:141–4.
65. Luu SU, Wang MF, Lin DL, et al. Ethanol and acetaldehyde metabolism in chinese with different aldehyde dehydrogenase-2 genotypes. *Proc Natl Sci Coun Repub China B*. 1995;19:129–36.
66. Thomasson HR, Crabb DW, Edenberg HJ, Li TK. Alcohol and aldehyde dehydrogenase polymorphisms and alcoholism. *Behav Genet*. 1993;23:131–6.
67. Wall TL. Genetic associations of alcohol and aldehyde dehydrogenase with alcohol dependence and their mechanisms of action. *Ther Drug Monit*. 2005;27:700–3.
68. Birley AJ, James MR, Dickson PA, et al. ADH single nucleotide polymorphism associations with alcohol metabolism in vivo. *Hum Mol Genet*. 2009;18:1533–42.
69. Birley AJ, James MR, Dickson PA, et al. Association of the gastric alcohol dehydrogenase gene ADH7 with variation in alcohol metabolism. *Hum Mol Genet*. 2008;17:179–89.
70. Eriksson CJ, Fukunaga T, Sarkola T, et al. Functional relevance of human adh polymorphism. *Alcohol Clin Exp Res*. 2001;25:157S–63.
71. Chen CC, Lu RB, Chen YC, et al. Interaction between the functional polymorphisms of the alcohol-metabolism genes in protection against alcoholism. *Am J Hum Genet*. 1999;65:795–807.
72. Chen WJ, Loh EW, Hsu YP, Chen CC, Yu JM, Cheng AT. Alcohol-metabolising genes and alcoholism among Taiwanese Han men: independent effect of ADH2, ADH3 and ALDH2. *Br J Psychiatry*. 1996;168:762–7.
73. Higuchi S, Matsushita S, Masaki T, et al. Influence of genetic variations of ethanol-metabolizing enzymes on phenotypes of alcohol-related disorders. *Ann N Y Acad Sci*. 2004;1025:472–80.
74. Higuchi S, Matsushita S, Muramatsu T, Murayama M, Hayashida M. Alcohol and aldehyde dehydrogenase genotypes and drinking behavior in Japanese. *Alcohol Clin Exp Res*. 1996;20:493–7.
75. Kimura M, Higuchi S. Genetics of alcohol dependence. *Psychiatry Clin Neurosci*. 2011;65:213–25.
76. Muramatsu T, Wang ZC, Fang YR, et al. Alcohol and aldehyde dehydrogenase genotypes and drinking behavior of Chinese living in Shanghai. *Hum Genet*. 1995;96:151–4.
77. Dickson PA, James MR, Heath AC, et al. Effects of variation at the ALDH2 locus on alcohol metabolism, sensitivity, consumption, and dependence in Europeans. *Alcohol Clin Exp Res*. 2006;30:1093–100.

78. Macgregor S, Lind PA, Bucholz KK, et al. Associations of ADH and ALDH2 gene variation with self report alcohol reactions, consumption and dependence: an integrated analysis. *Hum Mol Genet.* 2009;18:580–93.
79. Whitfield JB. Meta-analysis of the effects of alcohol dehydrogenase genotype on alcohol dependence and alcoholic liver disease. *Alcohol Alcohol.* 1997;32:613–9.
80. Whitfield JB, Nightingale BN, Bucholz KK, Madden PA, Heath AC, Martin NG. ADH genotypes and alcohol use and dependence in Europeans. *Alcohol Clin Exp Res.* 1998;22:1463–9.
81. Neumark YD, Friedlander Y, Thomasson HR, Li TK. Association of the ADH2*2 allele with reduced ethanol consumption in Jewish men in Israel: a pilot study. *J Stud Alcohol.* 1998;59:133–9.
82. Hasin D, Aharonovich E, Liu X, et al. Alcohol dependence symptoms and alcohol dehydrogenase 2 polymorphism: Israeli Ashkenazis, Sephardics, and recent Russian immigrants. *Alcohol Clin Exp Res.* 2002;26:1315–21.
83. Thomasson HR, Crabb DW, Edenberg HJ, et al. Low frequency of the ADH2*2 allele among Atayal natives of Taiwan with alcohol use disorders. *Alcohol Clin Exp Res.* 1994;18:640–3.
84. Wall TL, Carr LG, Ehlers CL. Protective association of genetic variation in alcohol dehydrogenase with alcohol dependence in Native American Mission Indians. *Am J Psychiatry.* 2003;160:41–6.
85. Gizer IR, Edenberg HJ, Gilder DA, Wilhelmsen KC, Ehlers CL. Association of alcohol dehydrogenase genes with alcohol-related phenotypes in a Native American community sample. *Alcohol Clin Exp Res.* 2011;35:2008–18.
86. McCarver DG. ADH2 and CYP2E1 genetic polymorphisms: risk factors for alcohol-related birth defects. *Drug Metab Dispos.* 2001;29:562–5.
87. McCarver DG, Thomasson HR, Martier SS, Sokol RJ, Li T. Alcohol dehydrogenase-2*3 allele protects against alcohol-related birth defects among African Americans. *J Pharmacol Exp Ther.* 1997;283:1095–101.
88. Jacobson SW, Carr LG, Croxford J, Sokol RJ, Li TK, Jacobson JL. Protective effects of the alcohol dehydrogenase-ADH1B allele in children exposed to alcohol during pregnancy. *J Pediatr.* 2006;148:30–7.
89. Viljoen DL, Carr LG, Foroud TM, Brooke L, Ramsay M, Li TK. Alcohol dehydrogenase-2*2 allele is associated with decreased prevalence of fetal alcohol syndrome in the mixed-ancestry population of the Western Cape Province, South Africa. *Alcohol Clin Exp Res.* 2001;25:1719–22.
90. Harada S, Okubo T, Nakamura T, et al. A novel polymorphism (–357 G/A) of the ALDH2 gene: linkage disequilibrium and an association with alcoholism. *Alcohol Clin Exp Res.* 1999;23:958–62.
91. Edenberg HJ, Jerome RE, Li M. Polymorphism of the human alcohol dehydrogenase 4 (ADH4) promoter affects gene expression. *Pharmacogenetics.* 1999;9:25–30.
92. Ehlers CL, Spence JP, Wall TL, Gilder DA, Carr LG. Association of ALDH1 promoter polymorphisms with alcohol-related phenotypes in southwest California Indians. *Alcohol Clin Exp Res.* 2004;28:1481–6.
93. Spence JP, Liang T, Eriksson CJ, et al. Evaluation of aldehyde dehydrogenase 1 promoter polymorphisms identified in human populations. *Alcohol Clin Exp Res.* 2003;27:1389–94.
94. Chou WY, Stewart MJ, Carr LG, et al. An A/G polymorphism in the promoter of mitochondrial aldehyde dehydrogenase (ALDH2): effects of the sequence variant on transcription factor binding and promoter strength. *Alcohol Clin Exp Res.* 1999;23:963–8.
95. Martin NG, Perl J, Oakeshott JG, Gibson JB, Starmer GA, Wilks AV. A twin study of ethanol metabolism. *Behav Genet.* 1985;15:93–109.
96. Kopun M, Propping P. The kinetics of ethanol absorption and elimination in twins and supplementary repetitive experiments in singleton subjects. *Eur J Clin Pharmacol.* 1977;11:337–44.



<http://www.springer.com/978-1-62703-046-5>

Alcohol, Nutrition, and Health Consequences

Watson, R.R.; Preedy, V.R.; Zibadi, S. (Eds.)

2013, XXI, 578 p. 70 illus., 8 illus. in color., Hardcover

ISBN: 978-1-62703-046-5

A product of Humana Press